

EPHOU 98-009

July 1998

hep-th/9807062

Prepotentials, Bi-linear Forms on Periods and Enhanced Gauge Symmetries in Type-II Strings

Takahiro Masuda *

and

Hisao Suzuki[†]*Department of Physics,**Hokkaido University**Sapporo, Hokkaido 060 Japan*

Abstract

We construct a bi-linear form on the periods of Calabi-Yau spaces. These are used to obtain the prepotentials around conifold singularities in type-II strings compactified on Calabi-Yau space. The explicit construction of the bi-linear forms is achieved for the one-moduli models as well as two moduli models with K3-fibrations where the enhanced gauge symmetry is known to be observed at conifold locus. We also show how these bi-linear forms are related with the existence of flat coordinates. We list the resulting prepotentials in two moduli models around the conifold locus, which contains α' corrections of 4-D N=2 SUSY $SU(2)$ Yang-Mills theory as the stringy effect.

*e-mail address: masuda@particle.sci.hokudai.ac.jp

[†]e-mail address: hsuzuki@particle.sci.hokudai.ac.jp

1 Introduction

In recent years there have been much progress about non-perturbative aspect in superstring theories. One of the non-trivial aspects of the superstring compactification is that the type-II strings with Calabi-Yau compactifications are connected by conifold transitions[1, 2]. Physically these are interpreted as the transitions through black hole condensation[3, 4, 5, 6]. Later, it was shown that some of these transition can be understand as ordinary higgs transition. These are discovered as a consequence of heterotic-type-II duality[7] where K3 fibered Calabi-Yau manifolds are shown to be a potential candidate for the dual of heterotic string compactification on $K3 \times T^2$ [8, 9, 10]. There has been many non-trivial check of this duality at one-loop order in refs.[11, 12, 13, 14, 15, 16, 17, 18, 19, 20]. The non-perturbative check has been given by showing the relation to Seiberg-Witten theory[21, 22, 23, 24]. Afterward, the generic strategy to read off the gauge groups from toric diagrams has been discussed[25].

As a practical analysis of the prepotential of the models, several methods has been studied in large moduli limit where we can argue them from mirror conjecture originated in ref.[26]. Convincing the mirror hypothesis with a consideration of monodromies, a systematic analysis has been established[27, 28, 29, 30]. On the other hand as for the prepotentials around conifold locus, there seems to be no systematic approach for the evaluation. However, in the case of enhanced gauge symmetry, an analysis has been given in ref.[21] where they solve the solutions of Picard-Fuchs equation around the points of enhanced gauge symmetries and the basis are chosen in such a way that the basis recovers the one in Seiberg-Witten basis in the limit $\alpha' \rightarrow 0$. In other words, the duality conjecture can be employed for the determination of prepotentials around the conifold locus, just like mirror conjecture have been in large moduli limit. As for the solutions of Picard-Fuchs equation, it is known that these are solved in the form of hypergeometric functions[31, 32, 29, 30] in the large moduli region. However, these preferable characterization seems to be lost around conifold locus although it may persists in some special cases[33, 34]. Off course, it is not an essential problem of the system whether solutions can be solved in compact expression. The main problem of the approach is how

to determine periods from the solutions of Picard-Fuchs equation. The usual approach for this is the consideration of monodromy[26, 27, 28]. However, the analysis for the generic situations, we may need the consideration of monodromies or the analytic continuation from the large moduli limit, which is expected to be more difficult when we have many moduli parameters. Therefore, it is favorable to have a new tool to determine periods at various region of moduli parameters.

In this article, we will construct bi-linear symplectic form on the solution space of Picard-Fuchs equations. Namely, we are going to consider a symplectic map from the solutions of the equations to c-numbers. Because it is manifestly invariant under monodromy transformations, we can determine the candidates of the periods up to the overall normalization which may also be determined by a simple consideration of monodromy. These bi-linear form will turn out to be related to the existence of flat coordinates, which will be explicitly demonstrated for one and two moduli models. In two moduli models which are shown to have enhanced gauge symmetries, we will show that the choice of periods determined by duality are consistent with the analysis of bi-linear form of the models which implies the integrability of the prepotentials. We will also list the explicit form of the prepotential around the point of the enhanced gauge symmetries from which we may argue how the stringy corrections enter for $SU(2)$ Seiberg-Witten theories[35].

This article will be organized as follows.

In section 2, we will show how to obtain symplectic form on the solution space of Picard-Fuchs equation by using the example of 1-moduli models and the 2-moduli models of $K3$ fibered Calabi-Yau three-fold. In section 3, we will demonstrate that the existence of flat coordinates are related directly to the symplectic form in one and two moduli models. In section 4, we will consistently specify the dual pair of fields around the conifold point in one and two moduli models by using bi-linear form. We will also list the prepotential around the point of enhanced gauge symmetries in two moduli models. Last section is devoted to the conclusion and some discussion.

2 Intersection form on the solution space of the Picard-Fuchs equations

In this section, we investigate how to determine the symplectic form of the solution space of the Picard-Fuchs equations associated to type II string theory. The construction of the symplectic basis defined in Gauss-Manin system is apparent for integrability and invariance of monodromy transformation. Thus we discuss first in this system though this system is equivalent to the Picard-Fuchs system.

Consider mirror manifold $\bar{M}$ of Calabi-Yau 3-fold M . Canonical homology basis of $H_3(\bar{M}, Z)$ is $\{A^a, B_a\}$ ($a = 0, 1, \dots, h^{2,1}$) and dual cohomology basis $\{\alpha_a, \beta^a\}$ of $H^3(\bar{M}, Z)$ are given by

$$\int_{A^a} \alpha_b = \delta_b^a, \quad \int_{B_a} \beta^b = \delta_a^b. \quad (a, b = 0, 1, \dots, h^{2,1}) \quad (2.1)$$

The period integral $f_i(\mu_a)$ is defined by

$$f_i(\mu_a) = \int_{\gamma_i} \Omega(\mu_a), \quad (2.2)$$

where Ω is a holomorphic 3-form, γ_i is a homology cycle in $H_3(\bar{M}, Z)$ and μ_a ($a = 1, \dots, h^{2,1}$) are moduli parameters of complex structure deformation.

Now consider Gauss-Manin system[36];

$$\frac{\partial}{\partial \mu_a} \Pi_i = M_a \Pi_i, \quad (i = 1, \dots, 2h_{2,1} + 2) \quad (2.3)$$

where $\Pi_i = {}^t(f_i, \partial_{\mu_a} f_i, \dots)$ is $(2h^{2,1} + 2)$ -dimensional vector and M_a is a $(2h^{2,1} + 2) \times (2h^{2,1} + 2)$ matrix. This system reduces to the Picard-Fuchs system satisfied by f_i . Associated to Gauss-Manin system, we define the symplectic form C_{ij} , as

$$\langle f_i, f_j \rangle = \Pi_i^T A \Pi_j = C_{ij}, \quad A^T = -A, \quad (2.4)$$

where second condition insures anti-symmetric form $C_{ij} = -C_{ji}$. On requiring C_{ij} to be independent from moduli parameter, we have to solve the equation $\frac{\partial}{\partial \mu_a} \langle f_i, f_j \rangle = 0$, i.e.,

$$\frac{\partial}{\partial \mu_a} A + M_a^T A + A M_a = 0. \quad (2.5)$$

Once we find A , we get the intersection form C_{ij} which defines moduli-independent norm on solution space. Under these conditions, it is easy to check the integrability of this system by using (2.1) and (2.3). Furthermore this form is manifestly invariant under monodromy transformation. Therefore the intersection form have the information enough to construct the period vector canonically. Namely, if we label f_i as f_a^A, f_a^B obeying following combinations

$$\langle f_a^A, f_b^A \rangle = \langle f_a^B, f_b^B \rangle = 0, \quad \langle f_a^A, f_b^B \rangle = \delta_{a,b}, \quad (2.6)$$

and set f_a^A to be a solution for A^a cycle, then f_a^B corresponds to a solution along B_a cycle naturally. For the canonical period vector $\{z_a, \mathcal{F}^a\}$, we identify $f_a^A = z_a, f_b^B = \mathcal{F}^b$ from this property up to normalization which undertaken by $Sp(2h^{2,1} + 2, R)$ transformation, where $\{z_a, \mathcal{F}^a\}$ are defined by

$$z_a = \int_{A^a} \Omega, \quad \mathcal{F}^a = \int_{B^a} \Omega. \quad (2.7)$$

$$\langle z_a, \mathcal{F}^b \rangle = \delta_a^b, \quad \langle z_a, z_b \rangle = \langle \mathcal{F}^a, \mathcal{F}^b \rangle = 0, \quad (2.8)$$

In this article, we intend to calculate prepotential of type II string theory by fixing period vectors from the Picard-Fuchs equation directly. So let us to discuss how to do this in the Picard-Fuchs system. In order to clarify the symplectic structure on Picard-Fuchs equations, we will use the following notation on the space of differential operators. Introducing bi-linear operator

$$B \wedge D(f_1, f_2) = \frac{1}{2}(Bf_1 Df_2 - Df_1 Bf_2), \quad (2.9)$$

where B, D are any differential operators with respect to moduli parameters μ_a , we will write the symplectic form in the form.

$$C_{ij} = C(f_i, f_j) = \sum_{k,k'} A_{(k,k')} D^{(k)} \wedge D^{(k')} (f_i, f_j), \quad (2.10)$$

where $A^{(k,k')}$ are functions of μ_a and $D^{(k)}$ are the k -th order differential operators. In order to obtain C_{ij} we have to find the combination of operator $D^{(k)} \wedge D^{(k')}$ in such a way that we have

$$\partial_{\mu_a} C = 0, \quad (2.11)$$

using the ring of differential operators given by Picard-Fuchs equation. If we can do this, next we solve this equation for $A_{(k,k')}$. This is an alternative way to find C_{ij} in the Picard-Fuchs system. Off course there is no guarantee that we will be able to have simple solutions for the coefficients $A_{(k,k')}$. However, as an explicit evaluations of the solutions for a few moduli models, we will obtain rather simple form of the symplectic forms, which will be considered in the next section.

2.1 1-moduli models

Before turning to the main subject of this article, we are going to perform how to obtain C_{ij} with the models of Calabi-Yau three-fold constructed from the hypersurfaces in the toric varieties and Grassmannian with 1-moduli as the basic examples.

For the models constructed by hypersurfaces in toric variety[31], Picard-Fuchs equation in the large complex structure limit can be written as

$$Df = \left[\theta_x^4 - x(\theta_x + \lambda_1)(\theta_x + 1 - \lambda_1)(\theta_x + \lambda_2)(\theta_x + 1 - \lambda_2) \right] f = 0, \quad (2.12)$$

where $\theta_x = x \frac{\partial}{\partial x}$, x is a variable made of the moduli of these manifolds being $x \sim 0$ in this region, and λ_i ($i = 1, 2$) is a rational number associated to each manifold[37, 38]. For example, the famous quintic model P^4 [5] corresponds to a model with $\lambda_1 = \frac{1}{5}, \lambda_2 = \frac{2}{5}$.

Let us consider the symplectic form of the solution space of above Picard-Fuchs equation. Since the Picard-Fuchs equation is of forth order, we propose the intersection form C up to third order of θ_x in the form:

$$C = A_1 1 \wedge \theta_x^3 + A_2 \theta_x \wedge \theta_x^2 + A_3 1 \wedge \theta_x^2 + A_4 1 \wedge \theta_x, \quad (2.13)$$

where coefficients A_i ($i = 1, \dots, 4$) are some functions of x to be determined. We impose the condition

$$\theta_x C = 0, \quad (2.14)$$

in the ring $D = 0$, so that the intersection of the solution space $\langle f_1, f_2 \rangle = C(f_1, f_2)$ is a constant for any pair of solutions of Picard-Fuchs equation. Applying θ_x to (2.9) and using

(2.7) to descend the order of θ_x^4 , we derive equations which have to be obeyed by A_i . The solution of the equations turns out to be very simple. We find the bilinear form is given by

$$C = (1-x) \left\{ 1 \wedge \theta_x^3 - \theta_x \wedge \theta_x^2 \right\} - x \left\{ 1 \wedge \theta_x^2(f_1, f_2) + [\lambda_1(1-\lambda_1) + \lambda_2(1-\lambda_2)] 1 \wedge \theta_x \right\}. \quad (2.15)$$

It seems interesting that the Wronskian of the Picard-Fuchs equation can be obtained by simple manipulation of C as $C \wedge C \sim (1-x) 1 \wedge \theta_x \wedge \theta_x^2 \wedge \theta_x^3$. In other words, C can be understood as a square root of Wronskian.

Let us see how this works when we construct the basis. In large moduli limit $x \sim 0$, a power solution of the Picard-Fuchs equation (2.6) can be written as

$$W_0 = \sum_{n=0}^{\infty} \frac{(\lambda_1)_n (1-\lambda_1)_n (\lambda_2)_n (1-\lambda_2)_n}{\Gamma(n+1)^4} x^n, \quad (2.16)$$

where $(a)_n = \Gamma(a+n)/\Gamma(a)$. Other three solutions, W_1, W_2, W_3 , can be given by Frobenius method; define the operator D_{ρ_i} applying to generic power solution $\sum_{n_i} c(n_i) x^{n_i}$,

$$D_{\rho_i} c(n_i) x^{n_i} = \left. \frac{\partial_{\rho_i} c(n_i + \rho_i)}{2\pi i} x^{n_i + \rho_i} \right|_{\rho_i=0}. \quad (2.17)$$

other three solutions can be written by

$$W_1 = D_{\rho_x} W_0, \quad W_2 = (D_{\rho_x})^2 W_0, \quad W_3 = (D_{\rho_x})^3 W_0. \quad (2.18)$$

In the leading order, these solution symplectic form can be written as

$$W_0 \sim 1, \quad W_1 \sim \frac{\log x}{2\pi i}, \quad W_2 \sim \frac{(\log x)^2}{(2\pi i)^2}, \quad W_3 \sim \frac{(\log x)^3}{(2\pi i)^3}, \\ C \sim 1 \wedge \theta_x^3 - \theta_x \wedge \theta_x^2, \quad (2.19)$$

Substituting these solutions directly to C in various combination, it is easy to see that only $C(W_0, W_3)$ and $C(W_1, W_2)$ are not zero, and all other combination vanish in the lowest order. Even including higher order terms, the intersection of these solutions are not changed. Therefore it is natural to conclude that the symplectic bases taken by these solutions are

$$z^0 \sim W_0, \quad z^1 \sim W_1, \quad \mathcal{F}_0 \sim W_3, \quad \mathcal{F}_1 \sim W_2, \quad (2.20)$$

where we impose the asymptotic behavior of $t = z^1/z^0$ is $t \sim \log x/(2\pi i)$. This means that if we fix the component of α cycle, the intersection form selects proper β cycle automatically. The intersection form by itself determines only whether the intersection of two solutions is zero or not, and this does not fix the normalization for the symplectic basis. In order to fix the normalization, we just need the classical yukawa coupling. Then, the prepotential is given by definition as

$$\mathcal{F} = \frac{1}{2} \frac{z^a \mathcal{F}_a}{(z^0)^2}, \quad (2.21)$$

and inverting t as $x = x(t)$, and substituting this to $\mathcal{F}$, the prepotential in this region is written as the function of t , which is equivalent to the result of [29, 30, 39].

In this way, the normalized bi-linear form can be obtained in the form;

$$C = \frac{(2\pi i)^3}{\kappa_0} \left[(1-x) \left\{ 1 \wedge \theta_x^3 - \theta_x \wedge \theta_x^2 \right\} - x \left\{ 1 \wedge \theta_x^2(f_1, f_2) + [\lambda_1(1-\lambda_1) + \lambda_2(1-\lambda_2)] 1 \wedge \theta_x \right\} \right], \quad (2.22)$$

where κ_0 is the classical yukawa coupling and $\kappa_0 = 16 \sin^2 \lambda_1 \pi \sin^2 \lambda_2 \pi$ for simple series of models [37, 38]. This normalized form can be considered as the intersection form on periods.

Next we extend the discussion to models constructed from the hypersurfaces in the Grassmannian considered recently in refs.[40]. A fundamental period in the large moduli limit can be obtained as diagonal degeneration of the periods of complete intersection of projective space[40]. For the examples of one moduli models listed in ref.[40], Picard-Fuchs operator of order 4 can be written in the following form as

$$\theta_x^4 = \sum_{i=1}^r a_i x^i (\theta_x^2 + i \theta_x + \alpha_i) (\theta_x^2 + i \theta_x + \beta_i), \quad (2.23)$$

where a_i, α_i, β_i are rational number associated to each case, and this reduces to the toric variety with $r = 1$ complete intersection with $a_1 = 1$, $\alpha_1 = \lambda_1(1-\lambda_1)$, $\beta = \lambda_2(1-\lambda_2)$. Along the same line in the case of toric variety, we can give the bi-linear form

$$C = (1 - \sum_{i=1}^r a_i x^i) \left\{ \theta_x \wedge \theta_x^2 - 1 \wedge \theta_x^3 \right\} + \sum_{i=1}^r i a_i x^i 1 \wedge \theta_x^2 + \sum_{i=1}^r (\alpha_i + \beta_i) x^i 1 \wedge \theta_x. \quad (2.24)$$

It is straightforward to specify periods even in these models.

2.2 2-parameter models with $K3$ -fibration

In this subsection we deal with a series of 2-moduli models of type II string theory compactified on Calabi-Yau three-fold constructed by $K3$ fibrations. This series consist of four models [8] denoted as $P^{1,1,2,2,6}[12]$, $P^{1,1,2,2,2}[8]$, $P^{1,1,2,2,2,2}[6, 4]$, $P^{1,1,2,2,2,2,2}[4, 4, 4]$. In these series, moduli parameters ψ and ϕ are combined to x and y as

$$x = \frac{\phi}{\psi^{\frac{1}{\lambda}}}, \quad y = \frac{1}{\phi^2}, \quad (2.25)$$

where λ is $\frac{1}{6}$, $\frac{1}{4}$, $\frac{1}{3}$, $\frac{1}{2}$ for $P^{1,1,2,2,6}[12]$, $P^{1,1,2,2,2}[8]$, $P^{1,1,2,2,2,2}[6, 4]$, $P^{1,1,2,2,2,2,2}[4, 4, 4]$, respectively[34]. Their Picard-Fuchs equations can be given by following set of operators

$$D_1 = \theta_x^2(\theta_x - 2\theta_y) - x(\theta_x + \lambda)(\theta_x + \frac{1}{2})(\theta_x + 1 - \lambda), \quad (2.26)$$

$$D_2 = \theta_y^2 - \frac{1}{4}y(2\theta_y - \theta_x + 1)(2\theta_y - \theta_x), \quad (2.27)$$

or equivalently,

$$D_1 = (1 - x)\theta_x^3 - 2\theta_x^2\theta_y - \frac{3x}{2}\theta_x^2 - [\frac{1}{2} + \lambda(1 - \lambda)]x\theta_x - \frac{\lambda(1 - \lambda)}{2}x, \quad (2.28)$$

$$D_2 = (1 - y)\theta_y^2 + y\theta_x\theta_y - \frac{y}{4}\theta_x^2 - \frac{y}{2}\theta_y + \frac{y}{2}\theta_x, \quad (2.29)$$

Now let us consider the combination of bilinear operators to construct the intersection form close in the order of θ as fewer as possible. The first order is trivial; $1 \wedge \theta_x$ and $1 \wedge \theta_y$. In second order there are four possibility; $1 \wedge \theta_x^2$, $1 \wedge \theta_y^2$, $1 \wedge \theta_x\theta_y$, $\theta_y \wedge \theta_x$. However one of these are not independent and we eliminate $1 \wedge \theta_y^2$ by using (2.18). Similarly in third order there are ten possibility, however, four terms which contain θ_y^2 , such as $1 \wedge \theta_x\theta_y^2$, $\theta_x \wedge \theta_y^2$, $\theta_y \wedge \theta_y^2$ and $1 \wedge \theta_y^3$, are descended by (2.18), and a term which contains θ_y^3 are eliminated by (2.17). Therefore in the third order we have five independent operator. Since the Picard-Fuchs operators are at most of order three in this series, we expect the intersection form close up to third order to be

$$\begin{aligned} C = & A_1 1 \wedge \theta_x^2\theta_y + A_2 \theta_y \wedge \theta_x^2 + A_3 \theta_x \wedge \theta_x\theta_y + A_4 \theta_x \wedge \theta_x^2 + A_5 \theta_y \wedge \theta_x\theta_y \\ & + A_6 1 \wedge \theta_x^2 + A_7 1 \wedge \theta_x\theta_y + A_8 \theta_y \wedge \theta_x + A_9 1 \wedge \theta_x + A_{10} 1 \wedge \theta_y. \end{aligned} \quad (2.30)$$

In order to determine coefficients A_i ($i = 1, \dots, 10$), we impose following conditions

$$\theta_x C = \theta_y C = 0, \quad (2.31)$$

in the ring of differential operators $D_1 = D_2 = 0$.

These will results in the first order differential equations for the coefficients. Although the equations seems to be messy, due to these relations, we can solve A_i in a simple form to find

$$\begin{aligned} C = & \frac{\Delta}{2(1-x)} 1 \wedge \theta_x^2 \theta_y - \frac{(1-x)(1-y)}{2} \theta_y \wedge \theta_x^2 - \frac{(1-x)(1-y)}{2} \theta_x \wedge \theta_x \theta_y \\ & + \frac{(2x-1)y}{4} \theta_x \wedge \theta_x^2 + (1-y) \theta_y \wedge \theta_x \theta_y - \frac{yx(2x+1)}{8(1-x)} 1 \wedge \theta_x^2 \\ & - \frac{x(1-y)}{4} 1 \wedge \theta_x \theta_y + \frac{x(1-y)}{4} \theta_y \wedge \theta_x \\ & + \frac{xy(x^2 + (-1 + 2\lambda(1-\lambda))x - 2\lambda(1-\lambda))}{8(1-x)} 1 \wedge \theta_x - \frac{x(1-y)\lambda(1-\lambda)}{2} 1 \wedge \theta_y, \end{aligned} \quad (2.32)$$

where $\Delta = (1-x)^2 - yx^2$ is the discriminant of the theory.

Let us see how above bi-linear form recovers canonical symplectic base which have been obtained previously. In the region $x \sim 0$, $y \sim 0$, power solution can be given by

$$W_0 = \sum_{n,m} \frac{(\lambda)_m (\frac{1}{2})_m (1-\lambda)_m}{\Gamma(m-2n+1) (m!)^2 (n!)^2} x^m \left(\frac{y}{4}\right)^m, \quad (2.33)$$

and other five solution are obtained by using Frobenius methods as before. As long as considering whether the intersection is zero or not, the normalization of the solution does not matter. Thus from the indicial equation we take logarithmic solutions as

$$W_1 = D_{\rho_x} W_0, \quad W_2 = D_{\rho_y} W_0, \quad (2.34)$$

$$W_3 = D_{\rho_x} D_{\rho_x} W_0, \quad W_4 = (D_{\rho_x} D_{\rho_y} + D_{\rho_x}^2) W_0, \quad (2.35)$$

$$W_5 = \left(\frac{2}{3} D_{\rho_x}^3 + D_{\rho_x}^2 D_{\rho_y}\right) W_0 \quad (2.36)$$

Furthermore, it is enough to calculate in the leading order when we construct the basis of period vectors. All solutions in the leading order can be written

$$W_0 \sim 1, \quad W_1 \sim \log x, \quad W_2 \sim \log y, \quad (2.37)$$

$$W_3 \sim (\log x)^2, \quad W_4 \sim (\log x + \log y)(\log x), \quad (2.38)$$

$$W_5 \sim \frac{2}{3} (\log x)^3 + (\log x)^2 \log y. \quad (2.39)$$

Also we can write the intersection form effectively in the leading order in the following form

$$C \sim \frac{1}{2} 1 \wedge \theta_x^2 \theta_y - \frac{1}{2} \theta_y \wedge \theta_x^2 - \frac{1}{2} \theta_x \wedge \theta_x \theta_y. \quad (2.40)$$

Each constant can be calculated directly as

$$C(W_0, W_5) = 1, \quad C(W_1, W_4) = -1, \quad C(W_2, W_3) = -1 \quad (2.41)$$

and the other combinations vanish. Assuming that $t_1 \sim z^1/z^0 \sim \log x/(2\pi i)$ and $t_2 \sim z^2/z^0 \sim \log y/(2\pi i)$, we conclude to take following combination as the symplectic base,

$$z^0 \sim W_0, \quad z^1 \sim W_1, \quad z^2 \sim W_2, \quad (2.42)$$

$$\mathcal{F}_0 \sim W_5, \quad \mathcal{F}_1 \sim W_4, \quad \mathcal{F}_2 \sim W_3, \quad (2.43)$$

Thus we can recover the result obtained in refs.[27, 29, 30]. In the connection to the normalization, the prepotential and Yukawa coupling of this series of models can be given by

$$\mathcal{F} = -\frac{1}{3!} K_{ijk} t^i t^j t^k + \dots, \quad (2.44)$$

$$K_{111} = 2K_{211} = 8 \sin^2 \lambda \pi. \quad (2.45)$$

All solutions are written with the normalization of the convention of [29] by using Yukawa coupling in appendix B. However, as was mentioned, we now concentrate on deriving the prepotential in one region, it is not necessary to fix the over all constant of the intersection form. All we need is the combination of the solution with non-zero intersection, and the relative normalization of Yukawa coupling of the tree level part of the prepotential, and also proper asymptotic behavior of α cycle.

3 Flat coordinate condition and bi-linear intersection form

This section is motivated by the observation in terms of the flat coordinate given in refs.[41, 42]. In string theory, you can calculate exact Yukawa coupling constant including instanton

effect in the large radius limit due to the mirror symmetry [26, 29, 30]. Along this calculation, flat coordinates are regarded as the special affine coordinates made of α cycles [43]; these are used to fix the combination of solution of Picard-Fuchs equation, so as to establish the property of maximal unipotent monodromy [26, 29, 30]. Recently interesting connection between periods and flat coordinate are worked out in [41, 42]. In these analysis, mirror symmetry can be represented as isomorphic pair of quantum cohomology ring of different manifold. By changing the basis of Jacobian ring, Gauss-Manin system can be written to a form with flat connection in terms of special coordinates; flat coordinates. Consequently the specification of the periods from these conditions completely match the results with regard to the maximal unipotency of monodromy [26, 29]. Since this monodromy property insures symplectic structure of periods, above analysis have to do with our procedure of specification of periods based on bi-linear form in the large radius limit. Our aim in this section is to show that bi-linear form we have constructed are directly related to the condition of flat coordinates.

3.1 1-moduli models

In this subsection we consider Gauss-Manin system of the Calabi-Yau space constructed from toric variety of 1-moduli models discussed in Sect. 2.1. Following refs.[41, 42], we consider a set of first order differential equations:

$$\theta_x w = G w, \quad (3.1)$$

where w is a vector which consists of period integrals. We prefer to use θ_x rather $\frac{\partial}{\partial x}$ to have to do with the result of last section, and choose basis to take $w = {}^t(f_i, \theta_x f_i, \theta_x^2 f_i, \theta_x^3 f_i)$ and f_i is given by (2.2) in the case of 1-moduli models. The Picard-Fuchs operator can be rewritten in the ring $D = 0$ by making use of (2.12) as

$$\theta_x^4 = a_4 \theta_x^3 + a_3 \theta_x^2 + a_2 \theta_x + a_1, \quad (3.2)$$

where

$$a_1 = \frac{\lambda_1(1-\lambda_1)\lambda_2(1-\lambda_2)}{1-x}x, \quad a_2 = \frac{\lambda_1(1-\lambda_1) + \lambda_2(1-\lambda_2)}{1-x}x,$$

$$a_3 = \frac{\lambda_1(1 - \lambda_1) + \lambda_2(1 - \lambda_2)}{1 - x}x, \quad a_4 = \frac{2x}{1 - x}. \quad (3.3)$$

In order for the Gauss-Manin system (3.1) to reduce the Picard-Fuchs equation (2.12), we take the matrix G as

$$G = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ a_1 & a_2 & a_3 & a_4 \end{pmatrix} \quad (3.4)$$

Let us introduce the flat coordinates $t = t(x)$ and new periods $v(t)$, to rewrite the Gauss-Manin system for v . Flat coordinate conditions can be represented as couplings of new basis given by degree preserving transformation of deformed Jacobian ring, which is reduced to the Gauss-Manin system for v in the following form:

$$\theta_t v(t) = R v(t) = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & K_{ttt}(t) & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix} v(t). \quad (3.5)$$

where $K_{ttt}(t)$ is Yukawa coupling in this coordinate. New period v is related to original one by

$$w = M(x) v, \quad (3.6)$$

and it is known that transfer matrix $M(x)$ can be taken as a lower-triangle matrix

$$M = \begin{pmatrix} r_{11} & 0 & 0 & 0 \\ r_{21} & r_{22} & 0 & 0 \\ r_{31} & r_{32} & r_{33} & 0 \\ r_{41} & r_{42} & r_{43} & r_{44} \end{pmatrix}. \quad (3.7)$$

By substituting (3.5) to (3.1) and rewriting to the Gauss-Manin system for v , we derive the condition to determine the transfer matrix M ,

$$A := GM - \theta_x M - \theta_x t M R = 0. \quad (3.8)$$

From the analysis about conditions $A_{i1} = 0$ ($i = 1, \dots, 4$), r_{11} must be the solution of the Picard-Fuchs equation, so we set $r_{11} = g$. Also after some algebra, we see that once we know concrete form of r_{44} , every r_{ij} and Yukawa coupling K_{ttt} can be represented by using $\theta_x t$ and r_{11} . To accomplish this, using components $A_{23}, A_{33}, A_{34}, A_{44}$ we derive the differential equation satisfied by r_{44} , so that we can determine the form of r_{44} :

$$r_{44} = \frac{1}{(1-x)g}. \quad (3.9)$$

Thus substituting this to all components but A_{43} , we obtain the complete form of the transfer matrix M .

Now let us show how flat coordinate condition relate to the bi-linear intersection form. The only component which have not been used to obtain r_{ij} is $A_{43} = 0$, or

$$\begin{aligned} & 2\theta_x g \theta_x^2 t + 4\theta_x t \theta_x^2 g + g \theta_x^3 t - 2 \frac{(\theta_x g)^2}{g} \theta_x t \\ & - \frac{x}{1-x} \left\{ g \theta_x^2 t + (\lambda_1(1-\lambda_1) + \lambda_2(1-\lambda_2)) g \theta_x t - 2\theta_x t \theta_x g \right\} = 0. \end{aligned} \quad (3.10)$$

As was claimed in ref.[42], the role of this condition is to determine the flat coordinate $t(x)$. Here we assume $t = f/g$, where f is a solution of the Picard-Fuchs equation. Although t is not necessary a ratio of periods, this assumption will make clear the relation between choice of the flat coordinate and specification of periods. Indeed, the condition $A_{43} = 0$ can be rewritten as the bi-linear intersection form for f and g itself!

$$C(g, f) = 0. \quad (3.11)$$

This means that in the 1-moduli models, to search for the flat coordinate under the above assumption is equivalent to determine the periods as a combination of the solutions of the Picard-Fuchs equation. Thus in the case of analyzing mirror symmetry, we may always take the flat coordinate in the large radius limit as the ratio of independent α cycles. The bi-linear intersection form is a key to explain this consequence explicitly.

In the 2-moduli models, the situation become slightly different from above in that the flat coordinate condition does not necessary coincide the condition of vanishing intersection of each period, as we will see in the next subsection.

3.2 2-moduli models

In this subsection we investigate how the flat coordinate condition relate to the bi-linear intersection form in 2-moduli models discussed in Sect.2.2.

Hereafter we set the basis of the ring $D_1 = D_2 = 0$ in 2-moduli models of (2.26) and (2.27) as $\{L^{(1)}, L^{(2)}, L^{(3)}, L^{(4)}, L^{(5)}, L^{(6)}\} = \{1, \theta_x, \theta_y, \theta_x^2, \theta_x \theta_y, \theta_x^2 \theta_y\}$. In this case, the Gauss-Manin system which recovers (2.26) and (2.27) can be written by

$$\theta_x w = G w, \quad \theta_y w = H w, \quad (3.12)$$

where $w = {}^t(f_i, L^{(2)} f_i, L^{(3)} f_i, L^{(4)} f_i, L^{(5)} f_i, L^{(6)} f_i)$, and G and H are given by

$$G = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ a_1 & a_2 & a_3 & a_4 & a_5 & a_6 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ a'_1 & a'_2 & a'_3 & a'_4 & a'_5 & a'_6 \end{pmatrix}, \quad H = \begin{pmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ b_1 & b_2 & b_3 & b_4 & b_5 & b_6 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ b'_1 & b'_2 & b'_3 & b'_4 & b'_5 & b'_6 \\ b''_1 & b''_2 & b''_3 & b''_4 & b''_5 & b''_6 \end{pmatrix}, \quad (3.13)$$

here we denote a_i etc. as coefficients for following operators in the ring $D_1 = D_2 = 0$

$$\theta_x^3 = a_i L^{(i)}, \quad \theta_x^3 \theta_y = a'_i L^{(i)}, \quad (3.14)$$

$$\theta_y^2 = b_i L^{(i)}, \quad \theta_x \theta_y^2 = b'_i L^{(i)}, \quad \theta_x^2 \theta_y^2 = b''_i L^{(i)}, \quad (3.15)$$

and all coefficients $a_i, a'_i, b_i, b'_i, b''_i$ ($i = 1, \dots, 6$) are immediately derived from the ring $D_1 = D_2 = 0$ after some algebra, though we shall not list them here. By introducing flat coordinates $t = t(x, y), s = s(x, y)$, and new period v relating to original one w as

$$w = M v \quad (3.16)$$

we intend to rewrite the Gauss-Manin system for v in the following form:

$$\theta_t v = R_t v, \quad \theta_s v = R_s v, \quad (3.17)$$

where, as the reduction of the flat coordinate condition of new basis of deformed Jacobian ring[42], matrix R_t and R_s are given by

$$R_t = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & K_{tts} & K_{ttt} & 0 \\ 0 & 0 & 0 & K_{tss} & K_{tst} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad R_s = \begin{pmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & K_{sts} & K_{stt} & 0 \\ 0 & 0 & 0 & K_{sss} & K_{sst} & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}. \quad (3.18)$$

In 2-moduli models, it is known that M needs non-vanishing components additionally to be a lower-triangle form as

$$M = \begin{pmatrix} r_{11} & 0 & 0 & 0 & 0 & 0 \\ r_{21} & r_{22} & r_{23} & 0 & 0 & 0 \\ r_{31} & r_{32} & r_{33} & 0 & 0 & 0 \\ r_{41} & r_{42} & r_{43} & r_{44} & r_{45} & 0 \\ r_{51} & r_{52} & r_{53} & r_{54} & r_{55} & 0 \\ r_{61} & r_{62} & r_{63} & r_{64} & r_{65} & r_{66} \end{pmatrix}. \quad (3.19)$$

By substituting (3.15) to (3.11) and comparing to (3.16), the condition to determine $M(x, y)$ is derived:

$$A := GM - \theta_x M - \theta_x t MR_t - \theta_x s MR_s = 0, \quad (3.20)$$

$$B := HM - \theta_y M - \theta_y t MR_t - \theta_y s MR_s = 0. \quad (3.21)$$

After some algebra, we see immediately r_{11} must be the solution of the Picard-Fuchs equation (2.26) and (2.27), so we set $r_{11} = g$ as before. Also we see that, as was pointed out in the ref.[42], some components coincide as

$$A_{i4} = A_{i5}, \quad B_{i4} = B_{i5} \quad (i = 4, 5, 6). \quad (3.22)$$

In order to know all r_{ij} , we must obtain r_{66} . To do this we concentrate on two conditions, $A_{44} = 0, A_{54} = 0$. Explicit calculation shows that there are identical terms in both conditions

up to factors. Consequently $\theta_y t A_{44} - \theta_x t A_{54} = 0$ will be the differential equation satisfied by r_{66} :

$$\begin{aligned}
& (\theta_x t)^2 \left\{ -b'_6 \theta_x - \theta_y - b'_6 \frac{\theta_x g}{g} - \frac{\theta_y g}{g} + b''_6 - (\theta_x b'_6) \right\} r_{66} \\
& + \theta_x t \theta_y t \left\{ 3\theta_x + a_6 \theta_y + 3 \frac{\theta_x g}{g} + a_6 \frac{\theta_y g}{g} - a_4 - a_6 b''_6 + a'_6 \right\} r_{66} \\
& + (\theta_y t)^2 \left\{ -2a_6 \theta_x - 2a_6 \frac{\theta_x g}{g} + a_4 a_6 + a_6 a'_6 - (\theta_x a_6) \right\} r_{66} = 0.
\end{aligned} \tag{3.23}$$

By solving this equation, we obtain r_{66} :

$$r_{66} = \frac{1-x}{\Delta g}, \tag{3.24}$$

where $\Delta = (1-x)^2 - x^2 y$ is the discriminant. By substituting this to suitable components of the condition (3.19) and (3.20), we can know exact form of M and Yukawa couplings, though we will not list these results here. We will rather turn to solve the flat coordinate condition. There still remain following components which we have not use until now

$$\begin{aligned}
A_{44} &= 0, \quad A_{54} = 0, \quad A_{64} = 0, \\
B_{44} &= 0, \quad B_{54} = 0, \quad B_{64} = 0.
\end{aligned} \tag{3.25}$$

These should be used to determine the flat coordinates $t(x, y)$, $s(x, y)$. How many conditions are independent among them? As a matter of fact, careful calculations reveal that four components A_{44} , A_{54} , B_{44} , B_{54} mean same condition up to factor:

$$\begin{aligned}
Q &:= (\theta_x t \theta_y s - \theta_x s \theta_y t) \left(-\frac{x}{2(1-x)} + \frac{\theta_x g}{g} - a_6 \frac{\theta_y g}{g} \right) \\
&\quad - \theta_x^2 t (b'_6 \theta_x s - \theta_y s) + \theta_x^2 s (b'_6 \theta_x t - \theta_x t) \\
&\quad + \theta_x \theta_y t (\theta_x s - a_6 \theta_y s) - \theta_x \theta_y s (\theta_x t - a_6 \theta_y t) = 0.
\end{aligned} \tag{3.26}$$

Furthermore, after complicated manipulations, it turns out that rest components A_{64} and B_{64} are represented by using derivative and multiples of Q as

$$A_{64} = -\frac{(b'_6 \theta_x t - \theta_y t)}{\Delta} \theta_x Q + \frac{(\theta_x t - a_6 \theta_y t)}{\Delta} \theta_y Q + \{\dots\} Q = 0, \tag{3.27}$$

$$B_{64} = -\frac{\{(b_4 + b_5 b'_6) \theta_x t - b'_6 \theta_y t\}}{\Delta} \theta_x Q + \frac{(b'_6 \theta_x t - \theta_y t)}{\Delta} \theta_y Q + \{\dots\} Q = 0. \tag{3.28}$$

where we denote negligible multiples of Q as dots in the brace. Therefore essentially flat coordinate condition is just (3.26). The question we have to ask now is how (3.26) relate to the bi-linear intersection form. Here we assume the flat coordinates are the ratio of the Picard-Fuchs equation as

$$t = \frac{f_1}{g}, \quad s = \frac{f_2}{g}, \quad (3.29)$$

where f_1, f_2 and g are independent solutions to each other. Under this assumption, resulting expression of (3.26) will be completely anti-symmetric in terms of g, f_1, f_2 , and the condition (3.26) can be rewritten by using the bi-linear intersection form (2.32) as

$$g C(f_1, f_2) + f_1 C(f_2, g) + f_2(g, f_1) = 0. \quad (3.30)$$

Thus we conclude that, contract to 1-moduli models, a set of vanishing conditions of mutual intersection of periods is a mere sufficient condition of the flat coordinate condition. Consequently, the solution space of the flat coordinate is larger than that of symplectic periods even under this assumption. So if you desire, you may find the solutions which satisfy the condition (3.26) although their mutual intersection does not vanish. Anyway, (3.26) represents manifestly the reason why the usual choice of the flat coordinate, which is a ratio of independent periods, work well.

4 Determination of prepotential around the conifold locus

4.1 1-moduli models around the conifold locus

In this subsection we turn to evaluate the value of symplectic form around the conifold point $x = 1$ in one moduli models. Changing the variable of (2.6) as $y = 1 - x$ and analyzing the incial equation, we put four solutions around the conifold Y_i ($i = 0, 1, 2, 3$) can be given in lowest order as

$$Y_0 \sim 1, \quad Y_1 \sim y, \quad Y_2 \sim y \log y, \quad Y_3 \sim y^2. \quad (4.1)$$

In this region, these solutions are obtained iteratively. The intersection of these solutions can be calculated directly in the leading order and we see that $C(Y_0, Y_3)$ and $C(Y_1, Y_2)$ are not zero and other combinations become zero. If we add higher order terms obtained by iterations, the intersection form (2.16) is not changed up to the higher orders as much as possible by mathematica. Taking conventional assumption for the asymptotic behavior for α cycle, it is natural to choose the symplectic base as

$$\tilde{z}^0 \sim Y_0, \quad \tilde{z}^1 \sim Y_1, \quad (4.2)$$

$$\tilde{\mathcal{F}}_0 \sim Y_3, \quad \tilde{\mathcal{F}}_1 \sim Y_2, \quad (4.3)$$

up to normalization. Around the conifold point, we cannot kill a degree of freedom such that we can add $\tilde{\mathcal{F}}_0$ to $\tilde{z}^0$ like $\tilde{z}'_0 = \tilde{z}_0 + \beta \tilde{\mathcal{F}}_0$, since this manipulation does not change the intersection of the basis. However, in other region $\tilde{\mathcal{F}}$ may be transformed by monodromy transformation, so from the global consistency of moduli space β must be a integer number. For maximal simplicity we set $\beta = 0$. The normalization of the periods can be fixed by the behavior around conifold locus [3, 4, 6]:

$$\tilde{\mathcal{F}}_1 \sim \frac{1}{2\pi i} \tilde{z}^1 \log \tilde{z}^1. \quad (4.4)$$

Therefore, we can calculate the prepotential around conifold point once we can specify the periods of α cycle. This procedure will be described in detail in rather non-trivial example with 2-moduli models in the following subsection.

4.2 prepotential around the conifold locus of 2-moduli model

In this section, we determine the period vector to give the exact prepotential of a series of 2-moduli models around the conifold point. Conifold transition of the models are discussed in Ref.[44] and the enhanced gauge symmetries has been established in Refs.[21, 22].

First of all, we have to obtain the solution of the Picard-Fuchs equation around the conifold point. On Calabi-Yau three-fold obtained by K3 fibrations with 2-moduli, three kinds of the power solution around the conifold point can be given in a systematic way[34]:

$$Y_0 = \sum_{n_1, n_2, m} \frac{(\frac{\lambda}{2})_{n_1} (\frac{1-\lambda}{2})_{n_1} (\frac{\lambda}{2})_{n_2} (\frac{1-\lambda}{2})_{n_2} \Gamma(n_1 + n_2 + 1)}{(n_1 + n_2 - 2n)! (\frac{1}{2})_{n_1} (n_1)! (\frac{1}{2})_{n_2} (n_2)! n! n!} x_2^{n_1 + n_2} \left(\frac{x_1}{4} \right)^m, \quad (4.5)$$

$$Y_1 = \sum_{n_1, n_2, m} \frac{(\frac{\lambda}{2})_{n_1} (\frac{1-\lambda}{2})_{n_1} (\frac{1+\lambda}{2})_{n_2} (1 - \frac{\lambda}{2})_{n_2} \Gamma(n_1 + n_2 + \frac{3}{2})}{\Gamma(n_1 + n_2 - 2n + \frac{3}{2}) (\frac{1}{2})_{n_1} (n_1)! (\frac{1}{2})_{n_2} (n_2)! n! n!} x_2^{n_1 + n_2 + \frac{1}{2}} \left(\frac{x_1}{4}\right)^m, \quad (4.6)$$

$$Y_2 = \sum_{n_1, n_2, m} \frac{(\frac{\lambda}{2})_{n_1} (\frac{1-\lambda}{2})_{n_1} (\frac{\lambda}{2})_{n_1} (\frac{1-\lambda}{2})_{n_1} \Gamma(n_1 + n_2 + 2)}{(n_1 + n_2 - 2n + 1)! (\frac{1}{2})_{n_1} (n_1)! (\frac{1}{2})_{n_2} (n_2)! n! n!} x_2^{n_1 + n_2 + 1} \left(\frac{x_1}{4}\right)^m, \quad (4.7)$$

where $x_1 = x^2 y / (1 - x)^2$, $x_2 = 1 - x$. The other logarithmic solutions can be represented formally by barnes-type representations including well-poised series of type ${}_4F_3$ [34]. However we could not find any good formula for the explicit evaluation of the representation. Fortunately, one solution Y_4 which is dual of Y_1 , is given by ordinary manner as

$$Y_4 = D_{\rho_y} Y_0. \quad (4.8)$$

In order to get remaining two solutions dual to Y_1 and Y_3 , which contain $\log(x_1 x_2^2)$ terms, we have to solve the equation for the coefficient iteratively by setting the solutions in the form:

$$Y_3 = \log(x_1 x_2^2) Y_0 + \sum_{n, m} p(n, m) x_1^n x_2^m, \quad (4.9)$$

$$Y_5 = \log(x_1 x_2^2) Y_2 + \sum_{n, m} q(n, m) x_1^n x_2^m. \quad (4.10)$$

From the point of view of heterotic-type II string duality, an enhancement of $SU(2)$ gauge symmetry occurs at the conifold point $x = 1$ so that, with suitable redefinition of the field and taking particle limit and decoupling gravity, the prepotential reduces to $SU(2)$ Seiberg-Witten prepotential. Let us denote new coordinate of moduli as $S = t_2 = W_2/W_0$, $T = t_1 = W_1/W_0$ where S is heterotic dilaton, and near $y = 0$, $x = 1$, S and T behave as

$$y = e^{-S} + \dots, \quad T = \frac{i}{2 \sin \lambda \pi} + \dots. \quad (4.11)$$

Around $SU(2)$ enhancement point $T = i/(2 \sin \lambda \pi)$, we introduce new coordinate $\tilde{T}$ as

$$\tilde{T} = i \frac{T - \frac{i}{2 \sin \lambda \pi}}{T + \frac{i}{2 \sin \lambda \pi}}, \quad (4.12)$$

where such transformation is performed explicitly in appendix C. This choice is just a generalization of the observation made in ref.[21]. Using these coordinates, prepotential around the conifold point can be written as

$$\mathcal{F} = \frac{1}{2} S \tilde{T}^2 + \mathcal{F}_{1-loop} + \mathcal{F}_{nonpert}. \quad (4.13)$$

Putting fields as

$$\tilde{T} = \sqrt{\alpha'} a, \quad e^{-S} = \alpha'^2 \Lambda^4 \exp(-\hat{S}), \quad (4.14)$$

$$x_1 = \frac{x^2 y}{(1-x)^2} = \frac{1}{u^2}, \quad x_2 = 1 - x = \alpha' u, \quad (4.15)$$

and taking particle limit $\alpha' \rightarrow 0$, we can fix the period vectors $(1, S, \tilde{T}, 2F - S\partial_S F - \tilde{T}\partial_{\tilde{T}} F, \partial_S F, \partial_{\tilde{T}} F)$ completely by asymptotic form in the leading order, which satisfy the above constraint;

$$\mathcal{F} = \alpha' \mathcal{F}_{SW} + O(\alpha'^2), \quad (4.16)$$

where $\mathcal{F}_{SW}$ is a prepotential of Seiberg-Witten theory in the weak coupling region. The specification of dual pair of fields in these models has been given in ref.[21]. What we are going to show here is that the choice is consistent of the integrability of prepotentials by using the symplectic form constructed in the previous section. In terms of Y_i , this choice of basis can be written in the lowest order as

$$\tilde{z}^0 \sim 1, \quad \tilde{z}^1 \sim \tilde{T} \sim x_2^{\frac{1}{2}}, \quad \tilde{z}^2 \sim S \sim \log(x_1 x_2^2), \quad (4.17)$$

$$\tilde{\mathcal{F}}_0 \sim x_2 \log(x_1 x_2^2), \quad \tilde{\mathcal{F}}_1 \sim x_2^{\frac{1}{2}} \log x_1, \quad \tilde{\mathcal{F}}_2 \sim x_2. \quad (4.18)$$

Next let us calculate higher order terms beyond the Seiberg-Witten prepotential. Going to the higher order, there are some ambiguity since this system include several kinds of integer power solutions; for example, we can add Y_2 to Y_0 without changing leading behavior in this limit. That is, the combination of these solutions can not determined by asymptotic behavior itself around the conifold point. This ambiguity is usually killed by the integrability of the prepotential. Without carrying out tedious long calculation of this type we instead use the intersection form, since it is insured to be integrable by using the intersection form.

As was the case with 1-moduli, it is enough to consider solutions up to first few orders to calculate the intersection, because this is not changed if the order of calculation become higher. Using the intersection form (2.25) and changing the variable to x_1, x_2 , we can give the intersection form which does not vanish as

$$C(Y_0, Y_5) = \frac{1}{4}, \quad C(Y_1, Y_4) = -\frac{1}{8}, \quad C(Y_2, Y_3) = \frac{1}{4}, \quad (4.19)$$

even including higher order terms. From this form, we can deduce the combination of period vectors, however there are some ambiguity of linear combination of the solutions which does not change value of intersection form. Now assuming proper asymptotic behaviors of α cycle $\tilde{z}^0 \sim 1$, $\tilde{z}^1 \sim 0$, $\tilde{z}^2 \sim \log y$, we fix this ambiguity in the basis.. There still remains an ambiguities, such as adding Y_2 to Y_3 as $Y'_3 = Y_3 + \beta Y_2$. The replacement is also consistent with the low energy behavior. From the global consistency of moduli space as before, β may be set to be zero. However we have not been able to fix the value at this moment.

Thus we conclude the symplectic base around the conifold point to be

$$\tilde{z}^0 \sim Y_0, \tilde{z}^1 \sim Y_1, \tilde{z}^2 \sim Y_3, \quad (4.20)$$

$$\tilde{\mathcal{F}}_0 \sim Y_5, \tilde{\mathcal{F}}_1 \sim Y_4, \tilde{\mathcal{F}}_2 \sim Y_2. \quad (4.21)$$

Using the definition of moduli as $\tilde{T} \sim z^1/z^0$, $S \sim z^2/z^0$, and taking relative normalization of Y_1 and Y_4 as $Y_4/Y_1 = (1/\pi i) \log x_1 + \dots$ which obeys the physical constraint of realizing Seiberg-Witten theory[35] in the lowest order[21], the prepotential of the theory is written by

$$\begin{aligned} \mathcal{F} = & \frac{1}{2} S \tilde{T}^2 + \frac{1}{2} \tilde{T}^2 \log \tilde{T}^4 + \frac{1}{2} \tau_0 \tilde{T}^2 + \frac{1}{64} e^{-S} \tilde{T}^{-4} \cdot \tilde{T}^2 - \frac{5}{32468} (e^{-S} \tilde{T}^{-4})^2 \cdot \tilde{T}^2 + \dots \\ & + \frac{-16 + 4\lambda - 4\lambda^2}{18} \tilde{T}^4 - \frac{8 - 8\lambda + 8\lambda^2}{96} e^{-S} \tilde{T}^{-4} \cdot \tilde{T}^4 + \dots \\ & + \dots \quad (4.22) \\ \rightarrow & \alpha' \mathcal{F}_{SW} + \alpha'^2 \left\{ \frac{-16 + 4\lambda - 4\lambda^2}{18} - \frac{8 - 8\lambda + 8\lambda^2}{96} \frac{\Lambda^4 e^{-\hat{S}}}{a^4} \dots \right\} + \dots \end{aligned}$$

where higher order terms are listed in the appendix B.

The degree of freedom of adding $\tilde{\mathcal{F}}_2$ to z^2 is reflected to $\tilde{T}^4$ term in the prepotential. On the contrary, the coefficient of the term $\tilde{T}^{2n}$ ($n \geq 3$) does not contain β . This imply that if S independent terms of $\mathcal{F}$ excepting logarithmic term is denoted as $h(T)$, which corresponds to the perturbative part, $\partial_T^5 h(T)$ becomes true modular form. This is consistent with the result of the heterotic perturbation theory.

5 Conclusion

We have shown how to get the prepotential around the conifold point in type-II string theory compactified on various types of Calabi-Yau three fold up to 2-moduli. We have introduced the intersection form which determine the symplectic form of period vectors of the solution of Picard-Fuchs equation. In 2-moduli model, it is easy to give the the exact prepotential around the conifold point by using fixed periods, which is represented as the correction of the stringy effect from the Seiberg-Witten theory. Unfortunately we could not fix a parameter in prepotential around conifold locus even by using both duality conjecture and bi-linear forms. We may need full knowledge of analytic continuation, or direct instanton calculation on the heterotic side.

Apart from the analysis of Calabi-Yau threefolds, we can obtain the bi-linear forms on Calabi-Yau d-folds[45]. As a quick analysis of one moduli models[46, 47], we find that even dimensional Calabi-Yau has symmetric forms contrary to the symplectic form in odd dimensions. These are consistent from the fact that intersection form of d-dimensional hypersurfaces are symmetric in even dimensions.

The treatment discussed in this article can be extended similarly to the other string theories, whose gauge symmetry enhanced at conifold locus are known, such as 3-moduli models. Anyway the non-perturbative prepotential which is given by means of bi-linear forms around the conifold locus are not directly verified at the moment, since we do not know the formulation of direct calculation for the non-perturbative effect in the heterotic string theory. We hope that some technique of the heterotic sting theory will be improved to estimate the justification of such macroscopic calculations some day.

A Normalization of periods in two-moduli models

In large complex structure limit, the prepotential and Yukawa coupling of a series of models we considering can be given by

$$\mathcal{F} = -\frac{1}{3!}K_{ijk}t^it^jt^k + \dots, \quad (\text{A.1})$$

$$K_{111} = 2K_{211} = 2K_{121} = 2K_{112} = 8 \sin^2 \lambda \pi. \quad (\text{A.2})$$

Using Yukawa coupling and setting $\tilde{D}_{\rho_i} = \frac{1}{2\pi i} D_{\rho_i}$, logarithmic solutions and coordinates of moduli space t_i can be written by [29, 30]

$$W_1 = \tilde{D}_{\rho_x} W_0, \quad W_2 = \tilde{D}_{\rho_y} W_0, \quad (\text{A.3})$$

$$W_3 = -\frac{1}{2} K_{211} \tilde{D}_{\rho_x}^2 W_0, \quad W_4 = -\frac{1}{2} (2K_{121} \tilde{D}_{\rho_x} \tilde{D}_{\rho_y} + K_{111} \tilde{D}_{\rho_x}^2) W_0, \quad (\text{A.4})$$

$$W_5 = \frac{1}{3!} (K_{111} \tilde{D}_{\rho_x}^3 + 3K_{112} \tilde{D}_{\rho_x}^2 \tilde{D}_{\rho_y}) W_0, \quad (\text{A.5})$$

$$t_1 = \frac{W_1}{W_0}, \quad t_2 = \frac{W_2}{W_0}. \quad (\text{A.6})$$

With this normalization prescription, if we set the normalization of the intersection form as $\tilde{C} = (K_{112}/(2\pi i)^3)C$, the intersection which does not vanish become $\tilde{C}(W_0, W_5) = \tilde{C}(W_1, W_4) = \tilde{C}(W_2, W_3) = 1$.

If we use $\tilde{C}$ around the conifold point to evaluate the intersection with taking the normalization of the solution as

$$Y_0 = N_0(1 + \cdots), \quad Y_1 = N_1(x_2^{\frac{1}{2}} + \cdots), \quad Y_2 = N_2(x_2 + \cdots), \quad (\text{A.7})$$

$$Y_3 = N_3(\log(x_1 x_2^2 + \cdots)), \quad Y_4 = N_4(x_2^{\frac{1}{2}} \log(x_1) + \cdots), \quad Y_5 = N_5(x_2 \log(x_1 x_2^2) + \cdots), \quad (\text{A.8})$$

the condition $\tilde{C}(Y_0, Y_5) = \tilde{C}(Y_1, Y_4) = \tilde{C}(Y_3, Y_2) = 1$ can be written as

$$N_0 N_5 = \frac{4K_{112}}{(2\pi i)^3}, \quad N_1 N_4 = \frac{-8K_{112}}{(2\pi i)^3}, \quad N_3 N_2 = \frac{-4K_{112}}{(2\pi i)^3}. \quad (\text{A.9})$$

Notice that if we set heterotic dilaton as $S = t_2 = (1/2\pi i) \log y + \cdots$, which is equivalent $N_3/N_0 = 1/(2\pi i)$, then $SU(2)$ Seiberg-Witten solution naturally arise because of the relation $N_4/N_1 = 1/(\pi i)$.

B Prepotential around the point of enhanced gauge symmetries in two-moduli models

The prepotential around $SU(2)$ enhancement point is given by

$$\mathcal{F} = \frac{1}{2} S \tilde{T}^2 + \frac{\tilde{T}^2}{2} \log \tilde{T}^4 + \frac{1}{2} \tau_0 \tilde{T}^2 + h(\tilde{T}) + \mathcal{F}_{n.p.}, \quad (\text{B.10})$$

where $\tau_0 = \log 4 - 6$ is the bare coupling. Coefficients of T in $h(T)$ up to order T^{12} are given by

$\tilde{T}^4$	$\frac{1}{18}(9b - 4(4 - \lambda + \lambda^2))$
$\tilde{T}^6$	$-\frac{2}{675}(1 - 14\lambda + 48\lambda^2 - 68\lambda^3 + 34\lambda^4)$
$\tilde{T}^8$	$\frac{2(1-2\lambda)^2}{99225}(10 - 305\lambda + 969\lambda^2 - 1328\lambda^3 + 664\lambda^4)$
$\tilde{T}^{10}$	$\frac{-4}{5.893025}(21 - 1638\lambda + 17432\lambda^2 - 80480\lambda^3 + 209596\lambda^4 - 335180\lambda^5 + 331648\lambda^6 - 188504\lambda^7 + 47126\lambda^8)$
$\tilde{T}^{12}$	$\frac{2}{3.25725625}(-883471770 + 22522670\lambda + 2343887\lambda^2 - 52343680\lambda^3 + 20001415\lambda^4 - 631410\lambda^5 + 18364886\lambda^6 - 25083360\lambda^7 + 18173880\lambda^8 - 7935360\lambda^9 + 1587072\lambda^{10})$

We give here non-perturbative part up to 5-th order from perturbative part. Non-perturbative part $\mathcal{F}_{n.p.}$ is decompose as $\mathcal{F}_{n.p.} = \sum_i \mathcal{F}_{n.p.}^i \tilde{T}^{i+2}$. Setting $p = e^{-S\tilde{T}^{-4}}$, each coefficient of p in $\mathcal{F}_{n.p.}^i$ is given by

$\mathcal{F}_{n.p.}^0$

p	$-\frac{1}{64}$	p^4	$-\frac{1469}{4.1073741824}$
p^2	$-\frac{5}{2.16384}$	p^5	$-\frac{4471}{5.34359738368}$
p^3	$-\frac{3}{54288}$		

$\mathcal{F}_{n.p.}^2$

p	$\frac{-1}{96}(-3\beta + 8(1 - \lambda + \lambda^2))$
p^2	$\frac{-1}{2.4096}(-9\beta + 4(4 - \lambda + \lambda^2))$
p^3	$\frac{7}{3.393216}(9\beta - 4(4 - \lambda + \lambda^2))$
p^4	$-\frac{715}{4.402953184}(-9\beta + 4(4 - \lambda + \lambda^2))$

$\mathcal{F}_{n.p.}^4$

p	$\frac{1}{1440}(-45\beta^2 + 240\beta(1 - \lambda + \lambda^2) - 2(153 - 282\lambda + 404\lambda^2 - 244\lambda^3 + 122\lambda^4))$
p^2	$\frac{-1}{2.18423}(117\beta^2 - 24\beta(20 - 11\lambda + \lambda^2) + 501 - 582\lambda + 800\lambda^2 - 436\lambda^3 + 218\lambda^4)$
p^3	$\frac{-1}{6.393216}(585\beta^2 - 720\beta(3 - \lambda + \lambda^2) + 2014 - 1436\lambda + 1832\lambda^2 - 7 - 2\lambda^3 - 396\lambda^3 + 396\lambda^4)$

$\mathcal{F}_{n.p.}^6$

p	$\frac{1}{4 \cdot 18900}(1575\beta^3 - 12600\beta^2(1 - \lambda + \lambda^2) + 210\beta(153 - 282\lambda + 404\lambda^2 - 244\lambda^3 + 122\lambda^4) - 4(6590 - 16935\lambda + 30789\lambda^2 - 313274\lambda^3 + 24552\lambda^4 - 10698\lambda^5 + 3566\lambda^6))$
p^2	$\frac{-1}{967680}(-5355\beta^3 + 1260\beta^2(28 - 19\lambda + 19\lambda^2) - 21\beta(3757 - 5318\lambda + 7536\lambda^2 - 4436\lambda^3 + 2218\lambda^4) + 60278 - 134739\lambda + 253751\lambda^2 - 278240\lambda^3 + 239660\lambda^4 - 120648\lambda^5 + 40216\lambda^6)$

 $\mathcal{F}_{n.p.}^6$

p	$\frac{-1}{6 \cdot 529200}(33075\beta^4 - 352800\beta^3(1 - \lambda + \lambda^2) + 8820\beta^2(153 - 282\lambda + 404\lambda^2 - 244\lambda^3 + 122\lambda^4) - 336\beta(6530 - 16935\lambda + 30789\lambda^2 - 31274\lambda^3 + 24552\lambda^4 - 10698\lambda^5 + 3626\lambda^6) + 8(165627 - 532161\lambda + 1158022\lambda^2 - 1568356\lambda^3 + 1635352\lambda^4 - 1188258\lambda^5 + 674168\lambda^6 - 238356\lambda^7 + 59586\lambda^8))$
-----	--

C Analytic properties of periods in $K3$ and torus

In this appendix, let us discuss in detail about the analytic properties of solutions. A part of this appendix will overlap with some results given recently in ref.[48]. As was discussed before, the difficulty of the analysis around the conifold point is the lack of the knowledge of the analytic continuation. However for the sake of $K3$ fibration, we may be able to continue three of six solutions which become power function around the conifold point. To see this, let us see the solution with no logarithmic term of y around large radius in the limit $y \rightarrow 0$. These functions are generalized hypergeometric function ${}_3F_2$:

$$W_0 = {}_3F_2(\lambda, \frac{1}{2}, 1 - \lambda; 1, 1, x) = \sum_n \frac{(\lambda)_n (\frac{1}{2})_n (1 - \lambda)_n}{n!n!n!} x^n, \quad (C.11)$$

$$W_1 = D_{\rho_x} W_0, \quad W_3 = \frac{1}{2} K_{211} (D_{\rho_x})^2 W_0. \quad (C.12)$$

Some formula for the hypergeometric function make us possible to continue to the conifold point. First of all, we rewrite the solution as the product of two hypergeometric function[49,

51, 46]

$${}_3F_2(\lambda, \frac{1}{2}, 1 - \lambda; 1, 1, x) = ({}_2F_1(\frac{\lambda}{2}, \frac{1 - \lambda}{2}; 1; x))^2. \quad (\text{C.13})$$

Next we are going to use the usual analytic continuation formula . However at this stage, naive continuation gives wrong result because this process exceed the branch. To implement this, we use quadratic transformation to rewrite the argument

$${}_2F_1(\frac{\lambda}{2}, \frac{1 - \lambda}{2}; 1; x) = {}_2F_1(\lambda, 1 - \lambda; 1; \frac{1}{2} - \frac{1}{2}(1 - x)^{\frac{1}{2}}). \quad (\text{C.14})$$

The formula of the analytic continuation of this type is well known. Therefore it is capable to continue ${}_3F_2$ function by using the representation of the product of two hypergeometric function

$$W_0 = ({}_2F_1(\lambda, 1 - \lambda; 1; \frac{1}{2} - \frac{1}{2}(1 - x)^{\frac{1}{2}}))^2. \quad (\text{C.15})$$

Similarly we can rewrite (D.18) as

$$\begin{aligned} W_1 &= {}_2F_1(\lambda, 1 - \lambda; 1; \frac{1}{2} - \frac{1}{2}(1 - x)^{\frac{1}{2}}) \cdot D_{\rho_x} {}_2F_1(\lambda, 1 - \lambda; 1; \frac{1}{2} - \frac{1}{2}(1 - x)^{\frac{1}{2}}) \\ &= {}_2F_1(\lambda, 1 - \lambda; 1; \frac{1}{2} - \frac{1}{2}(1 - x)^{\frac{1}{2}}) \cdot \frac{-1}{2i \sin \lambda \pi} {}_2F_1(\lambda, 1 - \lambda; 1; \frac{1}{2} + \frac{1}{2}(1 - x)^{\frac{1}{2}}), \end{aligned} \quad (\text{C.16})$$

$$\begin{aligned} W_3 &= \frac{K_{211}}{2} \left\{ (D_{\rho_x} {}_2F_1(\lambda, 1 - \lambda; 1; \frac{1}{2} - \frac{1}{2}(1 - x)^{\frac{1}{2}}))^2 \right. \\ &\quad \left. + \frac{1}{4 \sin^2 \lambda \pi} ({}_2F_1(\lambda, 1 - \lambda; 1; \frac{1}{2} - \frac{1}{2}(1 - x)^{\frac{1}{2}}))^2 \right\} \\ &= ({}_2F_1(\lambda, 1 - \lambda; 1; \frac{1}{2} + \frac{1}{2}(1 - x)^{\frac{1}{2}}))^2 + ({}_2F_1(\lambda, 1 - \lambda; 1; \frac{1}{2} - \frac{1}{2}(1 - x)^{\frac{1}{2}}))^2. \end{aligned} \quad (\text{C.17})$$

Now we continue these solution to the conifold point. Define

$$g_1(x) = \frac{\Gamma(\frac{1}{2})}{\Gamma(\frac{\lambda}{2} + \frac{1}{2})\Gamma(1 - \frac{\lambda}{2})} {}_2F_1(\frac{\lambda}{2}, \frac{1 - \lambda}{2}; \frac{1}{2}; 1 - x), \quad (\text{C.18})$$

$$g_2(x) = \frac{\Gamma(-\frac{1}{2})}{\Gamma(\frac{\lambda}{2})\Gamma(\frac{1 - \lambda}{2})} (1 - x)^{\frac{1}{2}} {}_2F_1(\frac{\lambda + 1}{2}, 1 - \frac{\lambda}{2}; \frac{1}{2}; 1 - x), \quad (\text{C.19})$$

and using the analytic continuation formula

$${}_2F_1(\lambda, 1 - \lambda; 1; \frac{1}{2} \pm \frac{1}{2}(1 - x)^{\frac{1}{2}}) = g_1(x) \mp g_2(x), \quad (\text{C.20})$$

results of the analytic continuation can be written in the form

$$W_0 = (g_1(x) + g_2(x))^2, \quad (\text{C.21})$$

$$W_1 = \frac{-1}{2i \sin \lambda \pi} (g_1(x) - g_2(x))(g_1(x) + g_2(x)), \quad (\text{C.22})$$

$$W_3 = 2(g_1(x)^2 + g_2(x)^2). \quad (\text{C.23})$$

The moduli parameter which is defined as the ratio of the two period is transformed as follows,

$$T = \frac{W_1}{W_0} = \frac{i}{2 \sin \lambda \pi} \frac{g_1(x) - g_2(x)}{g_1(x) + g_2(x)}, \quad (\text{C.24})$$

where near the conifold point, $g_1 \sim 1$ and $g_2 \sim 0$. This result is interpreted to the location of the gauge enhancement point in the moduli space

$$T = \frac{i}{2 \sin \lambda \pi}, \quad (\text{C.25})$$

which is the fixed point of the discrete subgroup of $SL(2, Z)$ of each model, especially if $\lambda = \frac{1}{6}$, this point is a fixed point of the transformation

$$T' = -\frac{1}{T}. \quad (\text{C.26})$$

Converting (3,30) to the expression for g_1 and g_2

$$\frac{g_2(x)}{g_1(x)} = \frac{T - \frac{i}{2 \sin \lambda \pi}}{T + \frac{i}{2 \sin \lambda \pi}}, \quad (\text{C.27})$$

we see that the ratio $g_2/g_1 = \tilde{T}$ is to be taken as the moduli around the conifold point, which is just the redefinition of the field around the enhanced point of gauge symmetry.

Though it is difficult to handle the analytic continuation of the solution with $\log y$, without carrying out this, we are able to write down the symplectic transformation from the period in the large moduli limit to the one around the conifold point, by imposing this transformation must be symplectic. As is expected from the transformation law of T , this transformation does not reduce to $Sp(6; Z)$ from $Sp(6; C)$ in general. In the case with $\lambda = 1/6$, if weak coupling behavior of heterotic dilaton is set to be $S = 1/(4\pi i) \log y + \dots$

rather than the prescription in appendix A, and moduli T is set to absorb the factor i as $T' = iT$, this transformation can reduce to $Sp(6; Z)$.

As was pointed out in ref.[8], in the limit $y \rightarrow 0$ $K3$ moduli τ reduces to the moduli of a kind of torus, which is represented by

$$\tau = \frac{i}{2 \sin \lambda \pi} \frac{{}_2F_1(\lambda, 1 - \lambda; 1; 1 - z)}{{}_2F_1(\lambda, 1 - \lambda; 1; z)}, \quad (\text{C.28})$$

where $z = \frac{1}{2} - \frac{1}{2}(1 - x)^{\frac{1}{2}}$, which we call $z_{\frac{1}{\lambda}}$. This is a very similar form to usual moduli of the torus

$$\tau = i \frac{{}_2F_1(\frac{1}{2}, \frac{1}{2}; 1; 1 - z_e)}{{}_2F_1(\frac{1}{2}, \frac{1}{2}; 1; z_e)}, \quad (\text{C.29})$$

however because of the factor $1/2 \sin \lambda \pi$, the relation between $z_{\frac{1}{\lambda}}$ and z_e is complicated. The relation to the absolute invariants are given in refs.[50, 8, 51]. We will give the relation by using various transformation of hypergeometric functions given in refs.[49, 52]. These identities has been used in the case of Seiberg-Witten theory[53].

- $\lambda = \frac{1}{6}$;

On τ side, from quadratic transformation [49]

$${}_2F_1(\frac{1}{6}, \frac{5}{6}; 1; z_6) = {}_2F_1(\frac{1}{12}, \frac{5}{12}; 1; 4z_6(1 - z_6)). \quad (\text{C.30})$$

On τ_0 side, using quadratic and cubic transformation

$$\begin{aligned} {}_2F_1(\frac{1}{2}, \frac{1}{2}; 1, z_e) &= {}_2F_1(\frac{1}{4}, \frac{1}{4}; 1, 4z_e(1 - z_e)) \\ &= (1 - z_e + z_e^2)^{-\frac{1}{4}} {}_2F_1(\frac{1}{12}, \frac{5}{12}; 1; \frac{27z_e^2(1 - z_e)^2}{4(1 - z_e + z_e^2)^3}). \end{aligned} \quad (\text{C.31})$$

Thus the relation of z_e and z_6 is

$$4z_6(1 - z_6) = \frac{27z_e^2(1 - z_e)^2}{4(1 - z_e + z_e^2)^3} = \frac{1}{J}, \quad (\text{C.32})$$

where J is the absolute invariant[49].

- $\lambda = \frac{1}{4}$;

On τ_o side, using quadratic transformation

$${}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; z_e\right) = \left(1 - \frac{1}{2}z_e\right) {}_2F_1\left(\frac{1}{4}, \frac{3}{4}; 1; \frac{z_e^2}{(2 - z_e)^2}\right). \quad (\text{C.33})$$

Thus z_4 is related to z_e

$$z_4 = \frac{z_e^2}{(2 - z_e)^2}. \quad (\text{C.34})$$

- $\lambda = \frac{1}{2}$;

In this case, the kind of the function on both side are same initially. However this does not mean $z_e = z_2$ because of the factor $1/(2 \sin \lambda \pi)$. In this case it is difficult to compare to each other, so, on τ side, by using quadratic transformation

$${}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; z_2\right) = (1 - z_2) \left(1 + \frac{4z_2}{(1 - z_2)^2}\right)^{-\frac{1}{2}} {}_2F_1\left(\frac{1}{4}, \frac{3}{4}; 1; \frac{4z_2}{(1 + z_2)^2}\right), \quad (\text{C.35})$$

we give the relation to $\lambda = \frac{1}{4}$ case

$$z_4 = \frac{4z_2}{(1 + z_2)^2}. \quad (\text{C.36})$$

Substituting this to (C.34) and solving for z_2 we read the relation to z_e in $\lambda = \frac{1}{2}$ case as

$$z_2 = \frac{z_e^2}{(1 + (1 - z_e)^{\frac{1}{2}})^4}. \quad (\text{C.37})$$

- $\lambda = \frac{1}{3}$; There seems to be any simple relations to other variables. However, we can find the relation to the absolute invariants by the quartic transformation[52]:

$${}_2F_1\left(\frac{1}{3}, \frac{2}{3}, 1, z_3\right) = (1 + 8z_3)^{-\frac{1}{4}} {}_2F_1\left(\frac{1}{12}, \frac{5}{12}, 1, \frac{64z_3(1 - z_3)^3}{(1 + 8z_3)^3}\right), \quad (\text{C.38})$$

as

$$\frac{1}{J} = \frac{64z_3(1 - z_3)^3}{(1 + 8z_3)^3}, \quad (\text{C.39})$$

as given in [8, 51]

Unfortunately, we have not found the formula taking critical role in $\lambda = \frac{1}{3}$ case.

D Intersection form in K3 manifold

In this appendix, we discuss briefly the possibility for applying our method by means of bi-linear form to different dimensional Calabi-Yau manifold. It is easy to expect that notion of symplectic intersection form directly acting on the solution space of the Picard-Fuchs equation, can be extended to arbitrary odd dimensional Calabi-Yau manifold. Now we concentrate on the even dimensional case. In this case homology of the manifold consist of $2n$ -dimensional hypersurfaces, and its dual basis of cohomology are of $2n$ -form. Therefore the interchange of these element appears to be symmetric. Taking into account of this situation, in spite of anti-symmetric operator (2.9), we way well construct symmetric bi-linear form by using anti-commuting operator

$$\{B, C\}(f_1, f_2) = \frac{1}{2}(Bf_1 Cf_2 + Cf_1 Bf_2). \quad (\text{D.40})$$

To be concrete, we are going to construct bi-linear form in the case of $K3$ manifold explicitly. Generally, the Picard-Fuchs operator of $K3$ manifold can be written by

$$D = \theta_x^3 - x(\theta_x + \lambda)(\theta_x + \frac{1}{2})(\theta_x + 1 - \lambda). \quad (\text{D.41})$$

Symmetric nature of cycles in this manifold is observed from the relation given by [46]

$$\int_M \alpha \wedge \beta = \int_M \beta \wedge \alpha = 1, \quad \int_M \gamma \wedge \gamma = 2, \quad (\text{D.42})$$

where α, β, γ are 2-forms corresponding to independent homology cycles. Now following the discussion of Section 2.1, we set bi-linear form C close up to second order of θ by means of operator (D.40) as

$$C = A_1\{1, \theta_x^2\} + A_2\{\theta_x, \theta_x\} + A_3\{1, \theta_x\} + A_4\{1, 1\}. \quad (\text{D.43})$$

By requiring the condition that C have to be constant

$$\theta_x C = 0, \quad (\text{D.44})$$

we can find coefficients A_i in following forms

$$A_1 = -2(1 - x), \quad A_2 = 1 - x, \quad A_3 = x, \quad A_4 = x\lambda(1 - \lambda). \quad (\text{D.45})$$

Immediately we can show that this bi-linear form can recover the previous result obtained in ref.[46] in the large radius limit. As a consequence, in the case of $K3$ space, method by means of bi-linear form can be used for specifying the period from the solution of Picard-Fuchs equation.

Since it is easy to see that we can extend above analysis to the case of higher even dimension, we conclude that the notion of bi-linear form can be employed in arbitrary even dimensional Calabi-Yau manifold, by introducing anti-commuting operator due to the symmetric nature of even dimensional homology cycle.

References

- [1] M. Reid, Math. Ann. **278**,329 (1987).
- [2] P.Candelas, A.Dale, C.A.Lutken and R.Schimmrigk, Nucl. Phys. **B298**, 493(1988).
- [3] A. Strominger, Nucl. Phys. **B451**, 96(1995).
- [4] B.R. Greene, D.R. Morrison, and A.Strominger, Nucl. Phys. **B451**, 109 (1995).
- [5] T.M.Chiang, B.R.Greene, M.Gross and Y.Kanter, Nucl. Phys. **B**, Proc. Suppl. **46**, 82(1996).
- [6] B. R. Greene, “*String Theory on Calabi-Yau Manifolds*”, hep-th/9702155.
- [7] S.Kachru and C.Vafa, Nucl. Phys. **B450**, 69(1995).
- [8] A.Klemm, W.Lerche and P.Mayr, Phys. Lett. **B357**, 313(1995).
- [9] P.S.Aspinwall and J.Louis, Phys. Lett. **B387**, 735(1996).
- [10] P.S.Aspinwall, *K3 surfaces and string duality*,hep-th/9611137.
- [11] A.Ceresole, R. D’Auria, S.Ferrara, W.Lerche, and J.Louis, Int. J. Mod. Phys. **A8**, 79(1993).
- [12] V.Kaplunovsky, J.Louis and S.Theisen, Phys. Lett. **B357**, 71(1995).
- [13] J.A.Harvey and G.Moore, Nucl. Phys. **B463**(1996)315; Phys. Rev. **D57**, 2329 (1998).
- [14] C.Kokorelis, *The master equation for the prepotential*, hep-th/9802099.
- [15] G.L.Cardoso, G.Curio, D.Lust and T.Mohaupt, Phys. Lett. **B382**, 241(1996).
- [16] G.L. Cardoso, G.Curio and D.Lust, Nucl. Phys. **B491**, 147 (1997).
- [17] M.Henningson and G.Moore, Nucl. Phys. **B482**, 187(1996).
- [18] T. Kawai, Phys. Lett. **B397**, 51(1997); *K3 surfaces, Igusa cusp form and string theory*, hep-th/9710016; *String duality and enumeration of curves by Jacobi forms* hep-th/9804014.
- [19] G.L.Cardoso, Nucl. Phys. Proc. Suppl. **56B**, 94(1997).
- [20] G.Moore,*String duality, automorphic forms, and generalized Kac-Moody algebras* hep-th/9710198.
- [21] S.Kachru, A.Klemm, W.Lerche, P.Mayr and C.Vafa, Nucl. Phys. **B459**, 537 (1996).
- [22] A.Klemm, W.Lerche, P.Mayr, C.Vafa and N.Warner, Nucl. Phys. **B477** 746 (1996).

- [23] W.Lerche, *Introduction to Seiberg-Witten Theory and its Stringy Origin*, hep-th/9611190.
- [24] A.Klemm, *On the Geometry behind $N=2$ Supersymmetric Effective Actions in Four Dimensions*, hep-th/9705131.
- [25] P. Candelas, E.Perevalov, G.Rajesh, Nucl. Phys. **B507**, 445(1997).
- [26] P.Candelas, X.de la Ossa, P.Green and L.Parks, Nucl. Phys. **B358**, 21(1991).
- [27] P.Candelas, X.de la Ossa, A.Font, S.Kats, D.R.Morrison, Nucl. Phys. **B416**, 482(1994).
- [28] P.Candelas, A.Font, S.Kats, D.R.Morrison, Nuc. Phys. **B429**, 626(1994).
- [29] S.Hosono, A.Klemm, S.Theisen and S.T.Yau, Comm. Math. Phys. **167**, 301(1995).
- [30] S.Hosono, A.Klemm, S.Theisen and S.T.Yau, Nucl. Phys. **B433**, 501 (1995).
- [31] V.V. Batyrev, *Dual Polyhedra and Mirror Symmetry for Calabi-Yau Hypersurfaces in Toric Varieties*, alg-geom/9310003.
- [32] V.V.Batyrev and D. van Straten, Commun. Math. Phys. **168**, 493 (1995).
- [33] H.Suzuki, Int.J.Mod.Phys. **A12**, 5123(1997).
- [34] H.Suzuki, Mod.Phys.Lett. **A12**, 2847(1997).
- [35] N.Seiberg and E.Witten, Nucl. Phys. **B426**, 19 (1994); Nucl. Phys. **B431**, 484 (1994).
- [36] P.Griffiths, Ann. Math. **90**, 460 (1969).
- [37] A.Klemm and S.Theisen, Nucl.Phys. **B389**, 153 (1993).
- [38] A.Libgover and J.Teitelbaun, Duke Math. J.**69**(1),29, *Int. Math. Res. Notes*(1993).
- [39] S. Hosono, *GKZ Systems, Gröbner Fans and Moduli Spaces of Calabi-Yau Hypersurfaces*, alg-geom/9707003.
- [40] V.V.Batyrev, I.C.Fontaine, B.Kim and D. van Straten, *Conifold Transitions and Mirror Symmetry for Calabi-Yau Complete Intersections in Grassmannians*, alg-geom/9710022; *Mirror Symmetry and Toric Degenerations of Partial Flag Manifolds*, math/9803108.
- [41] S.Hosono and B.H.Lian, *GKZ Hypergeometric Systems and Applications to Mirror Symmetry*, hep-th/9602147.
- [42] M.Noguchi, Int.J.Mod.Phys. **A12**, 4973 (1997).
- [43] A. Strominger, Commun. Math. Phys. **133**, 163(1990).
- [44] M. Lynker and R.Schimmrigk, Nucl. Phys. **B484**, 562 (1997).

- [45] B. R. Greene, D.R. Morrison and M.R. Plesser, Commun.Math.Phys. **173**, 559 (1995).
- [46] M. Nagura and K. Sugiyama, Int.J.Mod.Phys.**A10**, 233 (1995).
- [47] K. Sugiyama, Int.J.Mod.Phys. **A11**, 229 (1996).
- [48] M.Billo, F.Denef, P. Fre, I.Pesando, W.Troost, A.van Proeyen and D. Zanon, *The rigid limit in special Kähler geometry*. hep-th/9803228.
- [49] see for example, A.Erdélyi *et al.*, “*Higher Transcendental Functions*”,(McGraw-Hill, New York) Vol.**1**.
- [50] A. Klemm, B.H. Lian, S.S.Roan and S.-T. Yau, *A Note on ODEs from Mirror Symmetry*, hep-th/9407192.
- [51] B.Lian and S.T.Yau, Commun.Math.Phys. **176**, 163 (1996); *Mirror Maps, Modular Relations and Hypergeometric Series I*, hep-th/9507151, *Mirror Maps, Modular Relations and Hypergeometric Series II*, hep-th/9507153.
- [52] E. Goursat, Ann. Sci. Ecole Norm. Sup.(2) **10**, 3 (1881).
- [53] T.Masuda and H.Suzuki, Int.J.Mod.Phys. **A12**, 3413(1997).